


## Chapter 11 Problem 33 †



### Given

$$m_w = 25 \text{ kg}$$

$$m_b = 15 \text{ kg}$$

$$l = 1.6 \text{ m}$$

$$\omega = \frac{10 \text{ rev}}{\text{min}} \left( \frac{2\pi \text{ rad}}{1 \text{ rev}} \right) \left( \frac{1 \text{ min}}{60 \text{ s}} \right) = 1.05 \text{ rad/s}$$

### Solution

Find the angular momentum of the spinning barbell.

Each of the weights on the barbell are a distance of  $l/2$  from the pivot point. Each of the weights have a moment of inertia of

$$I_w = m_w r^2 = m_w (l/2)^2 = m_w l^2 / 4$$

The bar of the barbell is spun about its center. From table 10.2 of the textbook the bar has a moment of inertia of

$$I_b = (l/12) m_b r^2$$

The total moment of inertia of the barbell is

$$I = 2(m_w l^2 / 4) + m_b r^2 / 12$$

The angular momentum is then

$$L = I \cdot \omega = (2m_w l^2 / 4 + m_b r^2 / 12) \omega$$

$$L = [2(25 \text{ kg})(1.6 \text{ m})^2 / 4 + (15 \text{ kg})(1.6 \text{ m})^2 / 12] (1.05 \text{ rad/s})$$

$$L = 37.0 \text{ kg} \cdot \text{m}^2 / \text{s}$$

---

†Problem from Essential University Physics, Wolfson