


Chapter 13 Problem 45 †



Given

$$t = 6279 \text{ cycles}$$

$$m = 9.2 \text{ g} = 9.2 \times 10^{-3} \text{ kg}$$

$$L = 17 \text{ cm} = 0.17 \text{ m}$$

Solution

Find the length of the lecture.

The pencil is a physical pendulum and has a natural frequency determined by the equation

$$\omega = \sqrt{\frac{mgl}{I}} \tag{1}$$

Assuming the pencil has a uniform distribution of mass, $l = L/2$. The moment of inertia of the pencil is that of a rod rotated about one end. From Table 10.2 the moment of inertia is

$$I = \frac{1}{3}mL^2$$

Substituting these relationships into equation 1 gives

$$\omega = \sqrt{\frac{mgl}{\frac{1}{3}mL^2}} = \sqrt{\frac{mgL/2}{\frac{1}{3}mL^2}} = \sqrt{\frac{3g}{2L}}$$

The time period is

$$T = \frac{2\pi}{\omega} = \frac{2\pi}{\sqrt{\frac{3g}{2L}}} = 2\pi\sqrt{\frac{2L}{3g}} = 2\pi\sqrt{\frac{2(0.17 \text{ m})}{3(9.8 \text{ m/s}^2)}}$$

$$T = 0.676 \text{ s}$$

Since there are 6279 time periods during this lecture, the lecture must be

$$t = (6279 \text{ cycles})(0.676 \text{ s/cycle}) = 4243 \text{ s}$$

$$t = 70.7 \text{ min}$$

†Problem from Essential University Physics, Wolfson