

Chapter 16 Problem 76 †

Given

$$T_c = -15\text{ }^\circ\text{C}$$

$$T_0 = 20\text{ }^\circ\text{C}$$

$$C = 6.5 \times 10^6\text{ J/K}$$

$$R = 6.67\text{ mK/W} = 0.00667\text{ K/W}$$

Solution

Find the time for the house to cool to $0\text{ }^\circ\text{C}$.

Equation 16.3 is

$$Q = mc\Delta T$$

The heat capacity given in the problem is a combination of the mass and the specific heat of the house. Therefore $C = mc$ and

$$Q = C\Delta T$$

Q is measured in Joules and we want to find the heat flow in watts. As a differential amount of heat, dQ , leaves the house, the temperature changes by a differential amount, dT . Therefore, the differential form of this equation is

$$dQ = CdT$$

If the change is occurring with respect to time, then the heat flow in watts is

$$\frac{dQ}{dt} = C \frac{dT}{dt}$$

Next take equation 16.5

$$H = -kA \frac{\Delta T}{\Delta x}$$

Thermal resistance is defined as $R = \Delta x/(kA)$. This equation becomes

$$H = -\frac{\Delta T}{R}$$

Replace the difference in temperature with the current temperature, T , and the outside temperature, T_c , and we have

$$H = -\frac{T - T_c}{R}$$

Remember that $H = dQ/dt$; therefore,

$$-\frac{T - T_c}{R} = H = \frac{dQ}{dt} = C \frac{dT}{dt}$$

or

$$-\frac{(T - T_c)}{CR} = \frac{dT}{dt} \tag{1}$$

†Problem from Essential University Physics, Wolfson

This is a first order differential equation and can be solved by u-substitution. Let

$$u = T - T_c$$

$$du = d(T - T_c) = dT$$

Substitute into equation 1

$$-\frac{u}{CR} = \frac{du}{dt}$$

Separate the variables gives

$$-\frac{dt}{CR} = \frac{du}{u}$$

Integrating both sides will introduce an arbitrary constant and the result is

$$-\frac{t}{CR} + D = \ln(u)$$

Back substitute for u

$$-\frac{t}{CR} + D = \ln(T - T_c) \tag{2}$$

Exponentiate each side

$$e^{-t/CR+D} = T - T_c$$

Simplify

$$e^{-t/CR} \cdot e^D = T - T_c$$

The temperature at any particular time is

$$T = T_c + Ge^{-t/CR}$$

where $G = e^D$. This is the solution to Newton's law of cooling. This looks similar to decay problems with a time constant of CR . At $t = 0$, $T = T_0$, then

$$T_0 = T_c + Ge^{-0/CR} = T_c + G$$

Solving for G gives

$$G = T_0 - T_c = 20^\circ C - (-15^\circ C) = 35^\circ C$$

Since $G = e^D$, then

$$\ln(G) = \ln(e^D) = D$$

Therefore,

$$D = \ln(35^\circ C)$$

Using equation 2 solve for time when $T = 0^\circ C$.

$$-\frac{t}{CR} = \ln(T - T_c) - D$$

$$t = -CR [\ln(T - T_c) - D]$$

Substitute in the appropriate values gives

$$t = -(6.5 \times 10^6 \text{ J/K})(0.00667 \text{ K/W}) [\ln(0^\circ\text{C} - (-15^\circ\text{C})) - \ln(35^\circ\text{C})]$$

$$t = -(6.5 \times 10^6 \text{ J/K})(0.00667 \text{ K/W}) [\ln(15^\circ\text{C}) - \ln(35^\circ\text{C})] \quad (3)$$

$$t = 36,700 \text{ s} = 10.2 \text{ hours}$$

Some of you are bothered by the fact that I took the logarithm of a number with units, and you should be. But look closely at the brackets on the right side of equation 3.

Using the law of logarithms we get

$$\ln(15^\circ\text{C}) - \ln(35^\circ\text{C}) = \ln\left(\frac{15^\circ\text{C}}{35^\circ\text{C}}\right) = \ln(0.429)$$

This number is unitless and when substituted into equation 3 gives

$$t = -(6.5 \times 10^6 \text{ J/K})(0.00667 \text{ K/W}) \ln(0.429) = 36,700 \text{ s}$$

which is the same answer.