
## Chapter 18 Problem 45 $^{\dagger}$



## Given

$$\begin{split} \gamma &= 1.67 \\ P_A &= 250 \; kPa = 2.5 \times 10^5 \; Pa \\ V_A &= 1.00 \; m^3 \\ V_B &= 3.00 \; m^3 \end{split}$$

## Solution a) Find the pressure at *B*.

The process from A to B is adiabatic. The relationship between pressure and volume is then

$$PV^{\gamma} = const.$$

The pressure and volume at A and B are then related by

$$P_A V_A^{\gamma} = P_B V_B^{\gamma}$$

Solving for the pressure at B gives

$$P_B = \frac{P_A V_A^{\gamma}}{V_B^{\gamma}} = P_A \left(\frac{V_A}{V_B}\right)^{\gamma}$$
$$P_B = (2.5 \times 10^5 \ Pa) \left(\frac{1.0 \ m^3}{3.0 \ m^3}\right)^{1.67} = 3.99 \times 10^4 \ Pa$$

b) Find the pressure at C.

The process from C to A is isothermal. There relationship between pressure and volume is then

$$PV = nRT = const.$$

The pressure and volume at C and A are then related by

$$P_C V_C = P_A V_A$$

Solving for the pressure at C gives

$$P_C = P_A\left(\frac{V_A}{V_C}\right) = (2.5 \times 10^5 \ Pa)\left(\frac{1.0 \ m^3}{3.0 \ m^3}\right) = 8.33 \times 10^4 \ Pa$$

<sup>&</sup>lt;sup>†</sup>Problem from Essential University Physics, Wolfson

c) Find the net work done on the gas.

The work done during the adiabatic process is

$$W = \frac{P_A V_A - P_B V_B}{\gamma - 1}$$
$$W = -\frac{(2.5 \times 10^5 \ Pa)(1.0 \ m^3) - (3.99 \times 10^4 \ Pa)(3.0 \ m^3)}{1.67 - 1}$$
$$W = -1.94 \times 10^5 \ J$$

The work done during the process B to C is 0 J since the volume does not change. The work done during the isothermal process is

$$W = -nRT \ln\left(\frac{V_f}{V_i}\right)$$

Since nRT is constant, we can replace it with  $P_AV_A$ .

$$W = -P_A V_A \ln\left(\frac{V_A}{V_C}\right) = (2.5 \times 10^5 \ Pa)(1.0 \ m^3) \ln\left(\frac{1.0 \ m^3}{3.0 \ m^3}\right)$$
$$W = 2.75 \times 10^5 \ J$$

The total work is the sum of the work for all three processes.

$$W = -1.94 \times 10^5 \ J + 0 \ J + 2.75 \times 10^5 \ J$$
$$W = 8.1 \times 10^4 \ J$$