

Chapter 37 Problem 18 †

Given

$$I = 1.75 \times 10^{-47} \text{ kg} \cdot \text{m}^2$$

Solution

Find the wavelength emitted going from $l = 5$ to $l = 4$.

The rotational energy goes as

$$E_{rot} = \frac{\hbar^2}{2I}l(l+1)$$

Going from $l = 5$ to $l = 4$ results in an energy change of

$$\Delta E = E_4 - E_5 = \frac{\hbar^2}{2I}4(4+1) - \frac{\hbar^2}{2I}5(5+1) = \frac{\hbar^2}{2I}20 - \frac{\hbar^2}{2I}30 = -\frac{\hbar^2}{2I}10$$

This loss of energy is an energy gain by the photon. The energy of a photon is given by

$$E = \frac{hc}{\lambda}$$

Set this equal to the energy lost by the molecule gives

$$\frac{\hbar^2}{2I}10 = \frac{hc}{\lambda}$$

Replace \hbar with $h/2\pi$ and solving for wavelength gives

$$\frac{h^2}{8\pi^2I}10 = \frac{hc}{\lambda}$$

$$\lambda = \frac{hc8\pi^2I}{10h^2} = \frac{4\pi^2cI}{5h}$$

Substitute in the appropriate values gives

$$\lambda = \frac{4\pi^2(3.00 \times 10^8 \text{ m/s})(1.75 \times 10^{-47} \text{ kg} \cdot \text{m}^2)}{5(6.63 \times 10^{-34} \text{ J} \cdot \text{s})} = 6.25 \times 10^{-5} \text{ m} = 62.5 \text{ } \mu\text{m}$$

†Problem from Essential University Physics, Wolfson