Chapter 38 Problem 45 †

Given

 $m = 55.9206 \ u$ $m_p = 1.007276 \ u$ $m_n = 1.008665 \ u$ $1 \ u = 1.661 \times 10^{-27} \ kg$

Solution

Find the binding energy per nucleon for Iron-56.

Iron-56 has 26 protons and 30 neutrons. As free nucleons the total mass is

 $m_{free} = 26m_p + 30m_n = 26(1.007276\ u) + 30(1.008665\ u) = 56.4491\ u$

The mass difference between the free and bound nucleons is

$$\Delta m = m_{free} - m_{bound} = 56.4491 \ u - 55.9206 \ u = 0.5285 \ u$$

Converting this to kilograms gives

$$\Delta m = (0.5285 \ u) \left(\frac{1.661 \times 10^{-27} \ kg}{1 \ u}\right) = 8.778 \times 10^{-28} \ kg$$

The rest mass energy from special relativity is

 $E = mc^2$

$$E = (8.778 \times 10^{-28} \ kg)(3.00 \times 10^8 \ m/s)^2 = 7.90 \times 10^{-11} \ J$$

Convert the energy to MeV where $1 MeV = 1.6 \times 10^{-13} J$

$$E = (7.90 \times 10^{-11} J) \left(\frac{1 MeV}{1.6 \times 10^{-13} J}\right) = 494 MeV$$

Since there are 56 nucleons, then binding energy per nucleon is

 $\frac{494 \; MeV}{56 \; nucleon} = 8.82 \; MeV/nucleon$

[†]Problem from Essential University Physics, Wolfson